

## Methane Single Cell Protein: potential to secure a global protein supply against catastrophic food shocks

Juan B. García Martínez<sup>1\*</sup> , Joshua M. Pearce<sup>2</sup> , James Throup, Jacob Cates, Maximilian Lackner<sup>3,4</sup> , David C. Denkenberger<sup>1,5</sup> 

### Affiliations:

<sup>1</sup> Alliance to Feed the Earth in Disasters (ALLFED), Fairbanks, AK, USA.

<sup>2</sup> Department of Electrical & Computer Engineering, Western University, London, ON, Canada

<sup>3</sup> FH Technikum Wien, Höchstädtplatz 6, 1200 Wien, Austria

<sup>4</sup> Circe Biotechnologie GmbH, Brandstätte 1/27, 1010 Wien, Austria

<sup>5</sup> University of Alaska Fairbanks (Mechanical Engineering and Alaska Center for Energy and Power), Fairbanks, AK, USA 99775

\* Corresponding Author. Contact at: [juan@allfed.info](mailto:juan@allfed.info).

### **Abstract**

Global catastrophes such as a supervolcanic eruption, asteroid impact or nuclear winter could cause global agricultural collapse due to reduced sunlight reaching Earth's surface. Human civilization's food production system is unprepared to respond to such events, but methane single cell protein (SCP) could be a key part of the solution. Current preparedness centers around food stockpiling, an excessively expensive solution given that an abrupt sunlight reduction scenario (ASRS) could hamper conventional agriculture for 5 to 10 years. Instead, it is more cost-effective to consider resilient food production techniques requiring little to no sunlight.

This study analyses the potential of SCP produced from methane (natural gas and biogas) as a resilient food source for global catastrophic food shocks from ASRS. The following are quantified: global production potential of methane SCP, capital costs, material and energy requirements, ramp-up rates and retail prices. In addition, potential bottlenecks to fast deployment are considered.

While providing a more valuable, protein-rich product than alternatives, the production capacity could be slower to ramp up. Based on 24/7 construction of facilities, 7-11% of global protein requirements could be fulfilled at the end of the first year. Despite significant remaining

uncertainties, methane SCP shows significant potential to prevent global protein starvation during an ASRS at an affordable price — US\$3-5/kg dry.

## **Keywords**

Global Catastrophic Risk; Existential Risk; Single Cell Protein; Methanotrophic bacteria; Resilient food; Food Security; Global Catastrophic Food Shock; Nuclear Winter.

## **Highlights**

- Methane single cell protein has significant potential as resilient food for catastrophes.
- Protein requirements for the entire global population could be fulfilled in 2.5 to 4.5 years.
- Stranded natural gas (vented, flared or reinjected) could cover most of the requirements.
- The product would be affordable at an expected retail cost between US\$3-5/kg dry.
- Several recommendations and interventions were highlighted to expedite crisis response.

## **Abbreviations**

Abrupt sunlight reduction scenario (ASRS)

Capital expenditure (CAPEX)

Carbon dioxide (CO<sub>2</sub>)

Global catastrophic food shock (GCFS)

Natural gas (NG)

Net present value (NPV)

Single cell protein (SCP)

## 1 Introduction

The risk that a food shock will reduce global food production by about 10% has been estimated to be ~80% within this century (Bailey et al., 2015), and the chance of near-total food production loss could be as high as 10% (Denkenberger and Pearce, 2014, 2015). As the COVID-19 pandemic has shown, even disasters not directly related to the food system can substantially increase the population at risk of starvation (Laborde et al., 2020; Ahn and Norwood, 2020), and affect other major systems like energy (Brosemer et al., 2020), which are integral to the food system. Such events are most dire for countries critically dependent on food imports (Manero et al., 2020; Shokrani et al., 2020; Manheim and Denkenberger, 2020). A scenario on the order of magnitude of 10% food production loss, such as severe pollinator loss or abrupt climate change, could still result in mass starvation (Beasley, 2020). While “incremental” climate change from global warming over many decades is often considered in the food security literature, there has been limited discussion of more abrupt or extreme climate changes, which have the potential to create severe, sudden food shocks (Denkenberger and Pearce, 2015) with very short windows for adaptation.

There is a clear need for more work on preparedness. Several mechanisms exist for the occurrence of an abrupt global food shock due to agriculture’s dependency on stable environmental conditions such as sunlight, temperature and precipitation, as well as the dependency of agricultural yields on functioning supply chains. There is an increasing risk over time of concurrent severe weather events causing a multiple breadbasket failure which would exacerbate global food insecurity (Pham et al., 2022). However, the most extreme food shock that could threaten humanity in the near future is likely an *abrupt sunlight reduction scenario* (ASRS), in which a catastrophe causes a significant reduction in the amount of sunlight reaching the Earth’s surface. Potential causes include a “nuclear winter”, in which the atmosphere is clouded by soot from a global nuclear war causing the burning of cities, with less likely triggers including a supervolcanic eruption or asteroid/comet impact (Cirkovic, 2008; Denkenberger and Pearce, 2014). Consequences such as subzero temperatures across the Northern Hemisphere summer (Robock et al., 2007; Coupe et al., 2019) would preclude conventional agriculture for many years (Denkenberger et al., 2017), leading to a *global catastrophic food shock* (GCFS) from near-total global agricultural production loss. Such events, which would affect human wellbeing globally and even imperil modern civilization, are categorized as *global catastrophic risks* (Bostrom and Cirkovic, 2011). A severe ASRS is addressed in this study as an extreme limiting case scenario, in part because solutions that would work even under such circumstances could be useful for any scale of catastrophe. Global industry is considered to remain largely functioning after the catastrophe; Other GCFS scenarios involve a loss of industrial capacity (Denkenberger et al., 2021), requiring different solutions (Denkenberger et al., 2017). These types of events would demand radical innovation in food production, and a variety of complementary solutions would be required to forestall mass starvation: we named these *resilient food solutions* (Pham et al., 2022).

Given that an ASRS could last 5 to 10 years (i.e. nuclear winter), the cost of storing sufficient food for the global population is estimated to be extremely high in comparison to producing resilient foods that require less or no sunlight (Denkenberger and Pearce, 2015; Denkenberger et al., 2019). For example, in an ASRS cool-tolerant crops could be relocated to more adequate climates (Pham et al., 2022), simple greenhouses could be built on the tropics (Alvarado et al., 2020) and global seaweed production could be quickly ramped up (Mill et al., 2019), sugar could be produced from lignocellulosic biomass (Throup et al., 2022), synthetic fat could be produced from hydrocarbons

(García Martínez et al., 2022), acetic acid could be produced from CO<sub>2</sub> via microbial electrosynthesis (García Martínez et al., 2021a), mushrooms could be grown on residue from logging, cellulose-digesting ruminants and insects could be used as a food source (Denkenberger and Pearce, 2015), and leaf protein concentrate could be obtained (Pearce et al., 2019). This work studies the use of microbial protein produced via methanotrophic bacteria as a potential component of a food-crisis response. Resilient foods could be instrumental in avoiding starvation and sustaining society in the face of severe food shocks. Indeed, protein scarcity during wartime sparked initial interest in microbial protein (Ritala et al., 2017).

Microbial protein, referred to as single-cell protein (SCP), has been proposed as an alternative to animal products in meeting the growing global protein demand (Ritala et al., 2017). Outside of a disaster context, it is being considered as a sustainable alternative protein by an expanding body of literature (Claassens et al., 2016; Matassa et al., 2020; Leger et al., 2021; Khoshnevisan et al., 2022). A variety of microorganisms, including bacteria, fungi and microalgae, can be cultivated for SCP production (Ritala et al., 2017). Fungal SCP is already sold for human consumption under the brand name Quorn; however, the current production process relies on sugar (Ritala et al., 2017), a human-edible feedstock, making this product less useful in a GFCS. Conversely, during an extreme food supply crisis it could be desirable to produce SCPs for human consumption from non human edible feedstocks such as hydrocarbons, carbon dioxide (CO<sub>2</sub>) or available biowastes (Matassa et al., 2020), rather than using feedstocks such as sugars which could be more efficiently used directly for human consumption in such a dire situation. This would effectively increase food availability by virtue of obtaining a nutritionally rich product from inedible sources. Additionally, SCPs obtained from these resilient feedstocks would not compete on the input side with traditional protein sources. Methanol can also be used as an ASRS-resilient feedstock for SCP production, as was used for the first commercially available microbial protein product: Pruteen, from Imperial Chemical Industries. However, methanol is more localized and produced in smaller quantities than methane. Hydrocarbons can also serve as an ASRS-resilient feedstock for SCP production (Jenkins, 1988), specifically the paraffin components of petroleum which can also be used to produce synthetic fat (García Martínez et al., 2022).

Much research has focussed on SCPs on the grounds of sustainability due to their low water and land use (Matassa et al., 2015, 2016, 2020; Pikaar et al., 2018; Sillman et al., 2019). Methane-based bacterial SCP can use orders of magnitude less water and land than traditional protein sources such as meat and plant-based proteins. For example, Feedkind™ approximately uses 100 times less water and 1000 times less land per ton compared to soybeans (Cumberlege et al., 2016). Compared to other sunlight-independent food sources such as crops grown under artificial light or microalgae grown in photobioreactors, bacterial SCP has a very high energy efficiency (Alvarado et al., 2021). Due to its CO<sub>2</sub> emissions, SCP from natural gas is less sustainable than renewable hydrogen-based bacterial SCP, a similar type of SCP that could also be used as resilient food for GCFS (García Martínez et al., 2021b). However, natural gas SCP's lower requirements in terms of equipment imply a faster ramp-up and lower cost, which are fundamental characteristics for the potential of methane SCP as resilient food for GCFS (García Martínez et al., 2021b). This is because H<sub>2</sub> SCP requires construction of costly hydrogen production facilities while methane SCP can leverage existing natural gas extraction facilities. Most significantly for the purposes of crisis response, these types of SCP can produce high-quality, protein-rich food independently of traditional agriculture, specifically avoiding the need for sunlight and human edible inputs. For these closed-environment food production systems, exposure to several risk factors that affect conventional agriculture is greatly reduced or completely nullified, such as: institutional factors

(i.e. trade restrictions), abiotic factors (i.e. environmental degradation, extreme weather events, or climate variability), and biotic factors (i.e. pathogens or pests) (Tzachor et al., 2021).

Methanotrophic bacteria can be found in nature where methane naturally occurs, e.g. in swamps or tundra regions, and have been studied as a food source since the 1970s. Interest in them declined, but has been recovering thanks to their ability to produce high-quality protein with minimal land and water use. Methane SCP is one of the most advanced and accessible SCP production technologies, and is currently on the verge of large-scale commercialization (Flanagan, 2022). Methane is a relatively ubiquitous resource, and gas fermenters can be built at different scales, with a typical commercial plant size being 10,000 to 20,000 tons of protein per year. With several reactors being built on one site with joint use of utilities, economies of scale allow for production capacities on the order of 100,000 of SCP per year (Crumbley and Gonzalez, 2018; Flanagan, 2022).

Bacterial SCP from methane could potentially become an ingredient in a variety of food products, including solid foods like bread, pasta, and plant-based meats, as well as in liquid foods and drinks, such as plant-based dairy products, protein shakes, or broths (Southey, 2019). Bacterial SCP from methane has high protein completeness because its essential amino acid content is similar to, or higher than, the FAO guidelines (Ritala et al., 2017). Its amino acid content is superior to that of soybean meal, and it boasts a higher protein content in general (Pikaar et al., 2018). To date, there are no publicly available studies of bacterial SCP bioavailability in humans, but recent studies in fish point to bacterial SCP having a high digestibility (Glencross et al., 2020; Rajesh et al., 2022). Because of the fat content of methane SCP, its reported caloric content is 22 MJ/kg (Unibio Group, 2020a), noticeably higher than the average 17 MJ/kg caloric content of carbohydrates (U.S. Department of Agriculture, 2016). Methane SCP also boasts a notable micro-nutritional profile, containing significant amounts of B-group vitamins (esp. B1, B2, B3, B7, B8, B12), minerals (e.g. iron and magnesium) and essential fatty acids (Silverman, 2020; Duchene, 2016).

However, bacterial SCP has a high content of nucleic acids (8-12%) (Volova and Barashkov, 2010). This could cause health problems, such as gout and kidney stones (Ritala et al., 2017), if bacterial SCP were to be used as a significant food source for animals with long lifespans. Significant SCP consumption is not recommended for humans, unless the nucleic acid content is reduced during processing prior to use. The maximum safe limit of nucleic acid consumption for an adult human is 4 g/day (Adjei et al., 1995), which is equivalent to the amount present in 234 kcal worth of unprocessed bacterial SCP. This amount of unprocessed bacterial SCP in turn amounts to up to 59% of a daily recommended protein intake for human adults of 60 g/day. Unibio A/S claims to have developed a method to reduce nucleic acids to below 1% (Jorgensen, 2011).

Conversely, fungal SCP has been considered a safe component of the human diet for several years (U.S. Food and Drug Administration, 2002). Fungal SCP has a particularly low nucleic acid content compared to other SCPs. It is sold after post-processing that further reduces the nucleic acid content (Ritala et al., 2017) to below 2% of dry weight (Marlow Foods Ltd., 2001). If human adult daily protein requirements were to be fulfilled solely using fungal SCP, the total nucleic acid intake would still remain below the daily safe limit. Research has established that there are little to no threats associated with human consumption of fungal SCP, and this protein source is associated with a low incidence of allergic reactions compared to other sources (Finnigan et al., 2019). Although fungal SCP sets a favorable precedent, bacterial SCP by comparison is considerably less studied as a food source for humans, and requires more studies to establish its safety for human

consumption. However, according to Solar Foods, compositional data of their bacterial SCP product "raises no safety nor allergenicity concerns" (Solar Foods, 2021).

Key players currently pioneering mass production of methane SCP are Calysta Inc., Unibio A/S, Circe Biotechnologie GmbH, and String Bio Pvt Ltd. While these companies generally appear to be focusing on production of animal feed such as for the aquaculture sector, all four have explicitly shown interest in producing protein directly for human consumption (TEMASEK, 2020; Jensen, 2021; Lee, 2021; Circe Biotech, 2021). Separately, hydrogen-based SCP production companies such as Solar Foods and Air Protein are already developing the technology to produce human food-grade bacterial SCP and have applied for regulatory approval for human consumption (Solar Foods, 2021), which is a positive sign with regards both to the sustainability of the SCP for food concept but also for the possibility of consumer safety studies of bacterial SCP in the near future. This work assessed the viability of SCP as a direct human food source, rather than as an animal feedstock because the caloric conversion efficiency of animal products is low, at 3%-31% (Shepon et al., 2016), which is undesirable during a food-shortage scenario in which the intention is to maximize calories available for human consumption.

Two key metrics were used to globally characterize the potential of methane SCP as a resilient food: 1) how fast production can be ramped up to rapidly counter agricultural losses, and 2) how inexpensive the production cost is, as this determines affordability for the most economically disadvantaged populations. This work also addresses concerns about material constraints raised in previous studies of microbial protein as a resilient food in ASRS (Denkenberger and Pearce, 2014). The scope is akin to a FEL-1 stage (front-end loading) in which the concept is defined and preliminary budget estimates are produced, but the level of detail is not yet sufficient for construction (Warner, 2019).


The key aim is increasing resilience to global catastrophic risk, and more generally to existential risks with the potential to eliminate humanity or its future potential (Bostrom, 2013), since global famine could be considered an existential risk factor which could weaken our defenses to existential risk. In the face of these risks prevention is insufficient as a defense layer, response and resilience ought to be engaged as well, (Cotton-Barratt et al., 2020), which is the object of this work.

## 2 Methods

Methane SCP production requires three main inputs: 1) methane, which acts as both an electron donor and a carbon source, 2) a nitrogen source and 3) an oxygen source. Additionally, some minerals are also needed in smaller quantities. Anaerobic methane fermentation (Switzenbaum et al., 1990) has significantly lower yields and is hence not considered.

The chemical reaction used as a reference is the one proposed for *Methylococcus capsulatus* based on the premises proposed by (Villadsen et al., 2011), namely a yield coefficient of methane to cellular biomass of  $Y_{CH_4/biomass} = 0.8 \text{ g biomass/g CH}_4$  and a biomass degree of reduction  $k_x = 4.20$ ,

defined as the number of equivalents of available electrons per mol of carbon . The reaction can be expressed in a general form as shown in Equation 1.



Given ammonia as a nitrogen source for microbial growth, the resulting overall reaction can be expressed as shown in Equation 2. The formula  $CH_{1.8}O_{0.5}N_{0.2}$  stands for the simplified bacterial cell biomass, leaving aside sulfur and other minor components. The biomass is typically around 70% of protein and 30% of other compounds (e.g. fats, carbohydrates, ash).



The relevant unit operations, mass and energy flows involved in the proposed reference process are shown in Figure 1. The natural gas and  $O_2$  are fed together with ammonia and minerals to the bioreactor where the cell growth takes place in a continuous fermentation system, which gives a higher productivity than a batch fermenter. At the outlet of the reactor the fermentation broth contains an expected biomass concentration of 1-3% dry weight (i.e. bacterial cells), dissolved gasses and some unreacted ammonia and minerals. The water is removed via mechanical dewatering and drying steps, and the cells are disrupted in the homogenization step to increase digestibility by liberating the nutrients. The final SCP product is obtained in a powdered form, which may be subject to post-processing operations prior to storage. Water recycling, filters, pumps and heat exchangers are not shown.

A reduced nucleic acid content for the SCP product could be achieved via a number of processes, for example by applying a heat treatment (60-70 °C) to the effluent fermentation broth, or by using an alkaline treatment or chemical extraction (Ritala et al., 2017). Significant uncertainty remains as to which nucleic acid removal treatment would be employed in large-scale bacterial SCP production, but heat treatment is shown in Figure 1 in similarity to fungal SCP production. The heat treatment process activates endogenous, RNA-degrading enzymes for short time periods, while controlling for temperature and pH conditions (Anderson and Solomons, 1984). Once degraded, the nucleic acids diffuse out of the cell membrane and enter the liquid fraction, which has been previously separated from the SCP biomass through mechanical dewatering, for example through centrifugation. It may be possible to remove the need for heat treatments by instead using RNA-degrading enzymes at ambient conditions over longer periods of time, but this process has not been considered in the industrial production studied in this work. Different or no cell disruption techniques may be used instead of homogenization, but the lack of these may reduce digestibility.




Figure 1. Simplified process flow diagram of the reference methane SCP production process, based on (Jorgensen, 2011).

The process can use thermophilic bacteria to avoid excessive cooling; see e.g. (Levett et al., 2016) for a techno-economic assessment of the biopolymer PHB by methanotrophic bacteria. Pure or mixed culture operation has been described. The process contains an explosive atmosphere of methane and oxygen, meaning safety precautions have to be taken particularly in the headspace area. The methane source can be purified upstream of the reactor, or higher hydrocarbons can be consumed by a bacterial consortium.

## 2.1 Methodology overview

Two key metrics are estimated to characterize the potential of a resilient food for GCFS: the ramp up speed (how fast the production can be scaled over time) and the retail price per calorie (how affordable it would be during the catastrophe period). In addition, assessing the global availability of the relevant input resources is key to check for potential bottlenecks to fast production ramp-up. Figure 2 contains an overview of the methodology used to estimate these, which is described in depth in the following sections.




Figure 2. Methodology flowchart (TEA: techno-economic assessment, CAPEX: capital expenditure, OPEX: operating expenditure, NPV: net present value, CEPPI: Chemical Engineering Plant Cost Index).

## 2.2 CAPEX estimation

There is large uncertainty present in estimating the costs of a large scale "n-th plant" methane SCP production facility because currently no data on full-scale commercial size plants exist, "first of a kind" plants are costlier. The fixed capital expenditure (CAPEX) of the SCP plants was based on data from published industrial estimations by Unibio A/S. For a methane SCP production capacity of approximately 108,000 tonne dry product/year (tpa), the expected capital expenditure is \$251 million USD (Jorgensen, 2011). The cost estimate is hereby assumed to represent an n-th plant estimation, meaning that significant cost reductions after the technology reaches a higher level of maturity are not considered in the analysis. This production capacity is representative of that of currently projected full-scale commercial SCP production plants (Lane, 2018; Sefton, 2018; Rosenberry, 2019; Calysta Inc., 2020). After including nucleic acid removal from 9% (Unibio Group, 2020a) to 2%, the installed production capacity of the reference plant becomes 100,800 tpa. Preferably, many (5-10) bioreactors would be present rather than a single massive one for easier construction and operation given this large capacity; compare to the 1,500 m<sup>3</sup> airlift reactor used to produce Pruteen (Chisti, 1989).

The cost reference does not include the capital cost of O<sub>2</sub> production. This cost is estimated separately based on the methodology and values proposed by (Kreutz et al., 2005), including fixed and operating costs. The option of buying O<sub>2</sub> from industrial suppliers is discarded given the quantities involved in producing enough SCP to feed a significant portion of humanity. The updated capital expenditure of this air separation unit (ASU) is added to that of the Unibio reference factory to obtain the final plant CAPEX.

In an ASRS, food is expected to be scarce after the first few months, as food reserves run out (Denkenberger et al., 2017), making it preferable to increase food production as soon as possible. Fast construction methods are hereby proposed to reduce plant construction time, at the expense of increasing the capital expenditure. The fastest reasonable cost construction method available is

to implement 24/7 construction, reducing overall construction time to 32% of the original at an increased labor cost of 47% (Throup et al., 2022), according to the methodology and values of (Hanna et al., 2007). This value has been conservatively incorporated in terms of a 47% increase in the capital cost of the plant to account for labor constraints. State-of-the-art concurrent engineering could expedite the first steps of factory planning and engineering.

### **2.3 Assessment of required resources**

Natural gas (NG) is the largest source of methane available to industry today. One important alternative is biogas obtained from anaerobic digestion of organic matter, which can be used as a more sustainable alternative methane source (Cumberlege et al., 2016). If the nearly untapped global biogas potential was leveraged in its entirety, it could substitute the equivalent of 26-37% of the current natural gas production (Jain, 2019). Biogas has been proven as a feedstock for methane SCP production (Jones et al., 2020; Xu et al., 2021a), as well as electrochemically upgraded biogas (Acosta et al., 2020), and artificially synthesized methane from CO<sub>2</sub> (Xu et al., 2021b).

Different types of natural gas reserves can be defined (Attanasi and Freeman, 2013). Apart from economically exploitable reserves there are stranded gas reserves, which cannot currently be economically exploited for typical industrial uses. Physically stranded reserves cannot be accessed with current drilling technology. In contrast, economically stranded reserves are either 1) too far from their end use to justify transport costs, 2) contained in wells that are too small to justify extraction costs, or 3) associated with oil reserves and thus requiring extraction before the oil can be exploited, but not resulting in profit as a product. The last type is often regarded as an undesirable byproduct of oil extraction and is commonly flared, reinjected or vented. This associated natural gas is ripe for exploitation via methane SCP production because the methane is already being extracted without the need for additional capital cost. This is of considerable value in a GCFS because the resources that would be spent on extraction of further natural gas could instead be spent on the construction of more SCP plants or other resilient foods. In current times, exploiting stranded methane for production of SCP is estimated to be both economically feasible and more sustainable than the current status quo (El Abbadi et al., 2022). Currently a yearly average of nearly 150 billion cubic meters at standard temperature and pressure (bcm) of natural gas is being flared globally at upstream oil & gas and natural gas liquefaction plants (Elvidge et al., 2016; World Bank, 2018; EIA, 2019). An estimated yearly average of around 450 bcm of associated natural gas is being reinjected into oil wells (EIA, 2019).

Ammonia is considered as the nitrogen source due to widespread global availability from the fertilizer industry; another nitrogen source could be urea. Other potential nitrogen sources include gaseous ammonia recovered from biowaste pyrolysis or gasification (Matassa et al., 2015, 2020), as well as liquid digestate from anaerobic digestion (Khoshnevisan et al., 2019). During an ASRS, global agricultural production would most likely plummet, freeing up ammonia production capacity that could be used for SCP production instead. The oxygen is considered to be sourced from cryogenic air separation (Linde process), an industry standard process from which the majority of industrial oxygen is produced (FMI, 2019), as used by Calysta for methane SCP production (Cumberlege et al., 2016). It is recommended over other air separation processes (e.g. pressure swing adsorption or membrane separation) for a lower production cost in very large scale oxygen production (UIG, 2006; Alptekin, 2017). Using air directly as an oxygen source may be feasible but could significantly affect key performance parameters such as gas utilization, with major impact on the efficiency and economics of the process. Modeling and empirical studies,

outside the scope of this work, could be required to clarify this. Instead, this work considers the use of pure oxygen as feedstock in line with the practices of major methane SCP companies Calysta and Unibio (Cumberlege et al., 2016; Unibio Group, 2020b).

Resource requirements are summarized in Table 1, and are calculated based on estimates published for methane SCP, microbial protein requirements more generally, and for chemical industrial equipment. The required utilities to operate a bacterial SCP production plant of reference size are estimated in terms of electricity and fuel energy requirements. Due to the variability in properties of natural gas from different locations, a range of variables has been considered for the methane and energy content of the gas, which will result in a ranged estimation of the natural gas required to fulfill the food requirements. A ranged value of gas utilization has also been considered, which also broadens the overall range. A reactor designed for SCP production would have a gas utilization of at least 80% to be economical, while values of up to 90% have been reported in U-loop reactors (Jorgensen, 2011). The gas utilization affects both the natural gas and O<sub>2</sub> input requirements of the reference plant. Conservatively, the energy requirement for air separation has been selected as corresponds to high purity O<sub>2</sub>.

*Table 1. Basis of calculation for the energy requirements of methane SCP production.*

Variable	Value	Unit	Reference
Methane requirement	0.521	mol SCP/mol methane	(Villadsen et al., 2011)
Methane content of natural gas	87 - 98	%mole	(Enbridge Gas Inc., 2001)
Gas utilization of reactor	80 - 90	%	
Energy content of natural gas	35.4 - 42.8	MJ/m <sup>3</sup>	(Engineering ToolBox, 2005)
Energy content of SCP	22	MJ/kg	(Unibio Group, 2020a)
Solid content of dryer inlet	20	%	(Sillman et al., 2019)
Energy consumption of spray dryer	4880	kJ/kg evaporated water	(Baker and McKenzie, 2005)
Electricity to thermal energy usage ratio of spray dryer	1:27	Electricity:thermal	(Baker and McKenzie, 2005)
Electricity use of fermentation step	1.6	kWh/kg SCP	(Pikaar et al., 2018)
Electricity use of air	0.357	kWh/kg O <sub>2</sub>	(Aneke and Wang, 2015)

separation				
------------	--	--	--	--

Assessing potential bottlenecks to the ramp up potential of methane SCP first demands estimating the amount of SCP required to fulfill the food requirements of the global population and the equivalent in terms of the number of reference production plants. The amount of protein and calories available in the SCP product, as well as the requirements for feeding one person, were compared to the number of people globally. Then the resources required to produce the required amount of SCP are quantified, which includes energy and material resources such as natural gas as main feedstock, coal for the thermal requirements (if adequate), ammonia as a nitrogen source and electricity to run the process. The decision of whether to use coal or natural gas is not trivial and would depend on the specific location, since using natural gas may be more convenient but would reduce the amount available for SCP production if natural gas was the limiting factor in the location. Some of the natural gas that remains unused after passing through the reactor could be burned to produce energy, but this has been conservatively ignored. Additionally, alternative potential sources of methane are presented for discussion. The values used as a basis for the analysis are summarized in Table 2.

**Table 2.** Basis of calculation for the resource availability analysis. \*Some amount of food waste throughout the system is unavoidable, regardless of food crisis severity. However, a reasonably low value of food waste, 12%, was considered in the proposed scenario. This value was chosen because food waste is expected to be lower due to increased food scarcity. Moreover, the final bacteria SCP product is a dry product, with a long shelf life, further reducing potential food waste (Denkenberger and Pearce, 2014).

Variable	Value	Unit	Source
World population	7.8	billion people	(United Nations, 2019; Worldometers, 2020)
Recommended protein intake	60	g/person/day	(World Health Organization and United Nations University, 2007)
Expected food waste	12	% of calories produced	*
Average daily caloric requirement per person	2,100 (=1.39 kWh)	kcal/person/day	(World Health Organization, 2004)
Global ammonia production	171	Megatonne/year	(Research and Markets, 2020)
Global electricity consumption	2,551 (1 GWa = 8760, GWh = 8.76 TWh)	GW	(Sönnichsen, 2020)

Global coal production	7,337	Megatonne/year	(Rob Smith, 2018)
Global natural gas production	4,198	bcm/year	(IEA, 2019)
Global natural gas flaring and venting	150	bcm/year	(Elvidge et al., 2016; World Bank, 2018; EIA, 2019)
Global associated natural gas reinjecting	450	bcm/year	(EIA, 2019)
Global biogas production potential	26-37	% of current NG production	(Jain, 2019)

An accurate estimate of the protein content in the final SCP product is central to estimates of how much of it would be necessary to fulfill the protein requirements of the global population. A protein content range of 50-80% per kg of dry SCP product resulted from a review of multiple sources (Ravindra, 2000). This range was considered directly when determining, for the required SCP product amounts, the required resource amounts of the four possible energy sources: natural gas, electricity, coal, and ammonia.

## 2.4 Ramp-up speed estimation

We define the ramp-up speed as the increase in food production over time when continuously building as many food production factories as possible with the available resources. In the proposed catastrophe scenario, the ramp-up speed of methane SCP technology during a GCFS would likely be limited by the resources that could be effectively used, including but not limited to: raw materials, energy, qualified labor and capacity for equipment construction. We roughly account for these constraints by limiting the budget that can be effectively applied to 24/7 construction of methane SCP plants to a value of \$489 billion per year (Damodaran, 2020), which is the capital expenditure on adjacent industries whose resources could be redirected such as chemicals, power, pulp & paper, utilities and beverages. It is uncertain if workers of other, more disparate industries could be retrained fast enough to build and operate methane SCP plants. Thus, the average number of facilities that could be constructed in one year is obtained by dividing this total yearly CAPEX budget by the cost of a reference size factory.

The time taken to construct a facility is logarithmically related to the cost of the facility. The construction time was estimated by reference class forecasting, using a logarithmic regression model based on data from previously built factories (Martin et al., 2006). Given the urgency of the scenario, 24/7 construction is assumed to be used, which is estimated to reduce construction time to 32% of this value (Throup et al., 2022). The number of facilities that could be built per construction "wave" is calculated by dividing the amount of plants that can be built per year by the number of waves per year. For example, the first wave can be seen in the ramp up graphs as the first step increase in food production, shortly followed by another increase representing the moment that the factory transitions from startup production to full production, and later by another increase that represents the second wave.

The startup period is the time of reduced production between mechanical completion and the start of operation. An average production capacity of 50% applies, and it is considered to last one fourth of the construction time at regular speed (Humbird et al., 2011). Delays prior to plant construction also affect construction timelines. A value of 4 weeks is assumed, which is the time it took complex industries to convert and scale production of relevant supplies during the COVID-19 pandemic (Betti and Heinzmann, 2020). More details on ramp-up speed estimation can be found in (Throup et al., 2022) and (García Martínez et al., 2021b), including an example of the method on the supplementary material of the latter.

## 2.5 Economic analysis

A net present value (NPV) analysis was performed by calculating the required revenue for a standard unit of the SCP product when NPV equals zero. This analysis enabled estimates of the break-even cost of the product. To estimate the timeframe of plant operation, six years was used. This timeframe is shorter than those typical of chemical plants, and represents the period in which industrial food production factories could operate during an extreme food shock. This is representative of the duration for a period with little sunlight caused by a nuclear winter. The increased capital cost from 24/7 construction applies. At the end of the six year period the equipment was considered to be depreciated, corresponding to the time of coldest temperatures. In reality, some lower priced food could be sold for longer, there would be some salvage value, or the systems could be built less expensively (less durably), so this is a conservative assumption. To account for the time value of money, a 10% discount rate was used, consistent with recommendations for economic analyses facing an absence of statistical data for the given technology (Short et al., 1995). For comparison, the same analysis was performed for normal conditions outside of a catastrophe, namely a typical plant lifetime of 20 years and regular construction cost.

The operating costs are estimated based on the energy resource requirements for a reference plant from Table 4 and the prices from Table 3. For the electricity cost, the typical cost for the aluminum industry was taken to represent the low end of the uncertainty range, and the higher end of the uncertainty range was represented by the current European industry average. Thermal energy costs were calculated based on the cost of the amounts of coal required to reach thermal energy requirements. The total variable operating costs included the electricity and thermal energy costs described above, plus a further \$10.6 million USD for other operating costs, as well as a further 6.5\$ million USD for overheads (Jorgensen, 2011). Working capital was presumed at \$32.6 million USD. The financial parameters, with a federal tax rate of 35%, are based on the analysis of (Humbird et al., 2011). Financing sources were taken to comprise 70% equity (with an assumed 10% return on investment) and 30% loaned capital, assuming an interest rate of 8% per annum and a 10-year repayment period.

*Table 3. Energy resource cost ranges considered.*

Price range	Low	Middle	High
Electricity price	Global low (Burns, 2015)	U.S. average (EIA, 2020a)	Europe average (Eurostat, 2019)

(\$/kWh)		0.03	0.07	0.13
Natural gas price	Flared, vented or reinjected NG	10-year average (Markets Insider, 2020b)	10-year maximum (Markets Insider, 2020b)	
(\$/MWh NG)		0.00	11.26	16.51
Coal price	Global low (EIA, 2020b)	Average (Markets Insider, 2020a)	10-year high (Markets Insider, 2020a)	
(\$/tonne)		11.60	45.80	80.00

### 3 Results

#### 3.1 CAPEX estimation

The updated capital cost of the air separation unit required to fulfill the plant's O<sub>2</sub> requirements is estimated at \$67 million USD based on (Kreutz et al., 2005). Together with the other capital costs the total amounts to \$329 million USD for regular construction, or \$481 million for 24/7 construction. This would be equivalent to a CAPEX per unit of installed capacity of \$3,300/tpa and \$4,800/tpa, respectively. All costs are updated to 2020 values using the Chemical Engineering Plant Cost Index (CEPCI). These figures are conservatively estimated from the "first of a kind" plant proposed by Unibio; n-th plant facilities are at least 15% cheaper, in the authors' experience.

#### 3.2 Required resources and operating expenditures

The energy requirements estimated for each step are shown in Table 4. All values are calculated from Table 1: the natural gas requirement is obtained based on the proposed reaction stoichiometry, range of methane content, energy content of natural gas and gas utilization. The air separation energy requirement is calculated based on the energy use of the separation system from literature, the gas utilization and the reaction stoichiometry. The fermentation and centrifugation energy requirements are taken from a resource analysis study. Spray drying energy requirements are obtained from the industry average of a study on industrial spray drying data and the expected solid content of the inlet stream. The median energy requirements of spray

drying were lower, but the average was selected as a more conservative assumption. All values are corrected for nucleic acid removal.

*Table 4. Energy requirements of methane SCP production per step in kWh over dry mass of product.*

Step	Energy requirement (kWh/kg SCP)	
	90% gas utilization, low NG energy content	80% gas utilization, high NG energy content
Fermentation		1.6
Centrifugation		0.8
Spray drying		5.8
Air separation	1.5	1.7
Energy equivalent of required natural gas	21.2	32.5
Total energy requirements for methane SCP production	31.0	42.5

The energy analysis results for a reference plant are shown in Table 5. The values are estimated based on Table 4 and the production capacity. The calorific energy efficiency is estimated as the amount of energy invested in producing the SCP in comparison with its calorific content. The actual energy and efficiency values vary depending on the gas utilization and energy and methane contents of the gas. The estimated methane requirements have been compared against the value proposed by Bio Protein A/S (now part of Calysta). They stated a utilization of 2 m<sup>3</sup> of methane per kg of SCP produced (Babi and Price, 2010), in accordance with the value obtained here for high gas utilization, high methane content and no nucleic acid removal, which denotes the conservativeness of the present estimations.

*Table 5. Energy analysis results for a reference plant. The ranges are based on the intervals of gas utilization and energy content of the natural gas input considered.*

Variable	Low end	High end	Unit
Total energy requirements of reference plant	359	489	MW
Of which electricity is	48	50	MW

Overall energy efficiency		20.0	14.6	%
Natural gas requirement of SCP production		0.218	0.276	bcm/year
Thermal energy requirements in terms of:	Coal		62,039	tonne/year
	Natural gas	0.048	0.058	bcm/year


The share of global resources that would be required to fulfill the protein requirements of the global population via methane SCP is shown in Table 6 for both ends of the expected protein content range and gas utilization. The ammonia requirements are estimated based on the reaction stoichiometry. No resource bottlenecks are identified in comparison to the current availability.

Table 6. Range of the share of global resources required to fulfill the minimum global human protein requirements, while accounting for 12% food waste.

	<b>Low end</b>	<b>High end</b>	
Protein content of methane SCP	80	50	%
Gas utilization	80	90	%
Methane SCP requirement	243	388	Megatonne/year
Electricity capacity required	114	192	GW
Share of global electricity consumption	4.5	7.5	%
Natural gas required	524	1,062	bcm/year
Share of global natural gas production	12.5	25.3	%
Ammonia required		45	Megatonne/year
Share of global ammonia production		26.3	%
	Share of global natural gas production required	3.3	4.4 %
Thermal energy requirements in terms of:	Share of global coal production required	2.0	3.3 %

Even when accounting for fulfilling the entire global caloric requirements, the share of global natural gas that would have to be leveraged does not exceed 100%. At most it would be 97%, when assuming low gas utilization and use of natural gas for fulfilling the thermal requirements of the plants. The share of global electricity required in this case would be at most 25%. In comparison, the share of global ammonia production capacity required would be 110% of current values.

The amounts of natural gas required to fulfill global caloric and protein requirements via methane SCP are shown in Figure 3 in comparison to different available sources of methane. If current production of flared, vented and reinjected gas could be leveraged in its entirety for methane SCP production, it could cover 56-100% of the methane needed to fulfill global protein requirements, depending on protein content of the SCP, methane content of the gas and gas utilization. The nitrogen contained in the current production of sewage and manure is estimated between 3-13 megatonne/year (Matassa et al., 2020), equivalent to 8-36% of the nitrogen requirements for fulfilling the global protein requirements. However, it is unclear whether this could be leveraged for SCP production during a GCFS.



*Figure 3. Share of global caloric requirements that could be fulfilled by different potential methane sources (left) in comparison with the share of global protein requirements that could be fulfilled using the same sources (right).*

Bacterial SCP production also requires a number of minerals for cell growth metabolism. These include calcium, iron, phosphorus, sulfur, sodium, chlorine, potassium, and magnesium, among others. Out of these, magnesium was the only one identified as a potential bottleneck to SCP production ramp-up (García Martínez et al., 2021b), but only when aiming to fulfill global caloric requirements.

### 3.3 Ramp-up speed values

The construction time for a reference size plant is estimated at 87 weeks, and at 27 weeks when using 24/7 construction. The ramp up speed for the scenario in which the global budget for chemical and related industries can be effectively redirected to fast construction of methane SCP factories is shown in Figure 4 for the global caloric requirements and Figure 5 for the global protein requirements. For the fast construction scenario at the end of the first year around 2% of the caloric requirements could be fulfilled, translating to 7-11% of the protein requirements. The global protein requirements could potentially be covered in approximately 2.5-4.5 years. Note that the regular construction speed eventually overtakes 24/7 construction due to its lower resource intensity, but takes much longer to start producing food which makes it worse in the advent of a GCFS.





Figure 4. Expected ramp up speed of methane SCP production in terms of the global caloric human requirements fulfilled over time. The results shown reflect the use of the budget of similar industries, including regular and fast construction speeds.



*Figure 5. Expected ramp up speed of methane SCP production in terms of the global protein human requirements fulfilled over time, for different values of the protein content of the SCP product. The results shown reflect the use of the budget of similar industries, including regular and fast construction speeds.*

For reference, if assuming unlimited capital and no bottlenecks, a capital cost of 6.1 trillion USD would be sufficient for building the amount of methane SCP factories required to fulfill the caloric requirements of humanity. This would take an amount of time equivalent to that of building one reference scale production plant at around 13 months to full production (see the second step in the curves from Figures 3 and 4). Assuming slow construction, the capital cost would amount to 4.1 trillion USD, corresponding to a timeframe of 27 months until full production levels would be achieved.

Humans are unlikely to survive by relying on methane SCP as their sole food source; aiming to fulfill all global human caloric requirements through SCP alone is unrealistic. However, methane SCP could potentially fulfill the protein requirements of the global population over the duration of the proposed sunlight reduction scenario, approximately equivalent to 19-31% of caloric requirements. This amount of protein could be provided through SCP production on a construction budget of 1.2-1.9 trillion USD, a range that assumes fast construction methods and varies depending on the protein content per unit of SCP product. At a fast construction budget of around 1.2-1.9 trillion USD depending on protein content. The middle of the range of protein content, 65%, corresponds with the protein content expected from the Unibio SCP reference, assuming it has had nucleic acids removed, so it can be considered to constitute a "best estimate" of the ramp-up speed. Based on it, the protein requirements would be fulfilled in 3.5 years with the limited construction budget. A similar calculation based on a 9.8% fat content of the methane SCP

(Silverman, 2020) and a minimum recommended fat intake equivalent to 15% of total energy intake (García Martínez et al., 2022) yields about 25% coverage of the global fat requirement when fulfilling the entire global protein requirement via methane SCP in this way. Other resilient foods could cover the rest of the nutritional requirements (Pham et al., 2022).

### 3.4 Food price

The NPV analysis was performed to estimate the break-even cost of the methane SCP product for different scenarios. The expected cost of the SCP in the ASRS was estimated by limiting the plant life to 6 years and accounting for the additional cost of 24/7 construction. For comparison, the product cost in regular conditions (20 years of lifetime and regular construction cost) was also obtained. For each of the two scenarios, the product cost was calculated for a scenario of high operating costs (high natural gas and electricity cost), and for an optimistic scenario using free natural gas (i.e. vented or flared) and low electricity cost, as lower and upper bounds for the cost. Results are shown in Figure 6.




Figure 6. Breakdown of the contributions to the wholesale production cost incurred per unit of methane SCP produced.

A markup of 100% was applied to estimate the retail cost of the SCP product, accounting for distribution and other additional costs (McCray, 2010). We refer to these values as a retail cost

instead of price due to the uncertain equilibrium of the market during a GCFS, which could alter the sale price. The result is shown in Table 7. The retail cost for fulfilling a person's daily caloric requirements would be \$1.19-2.03.

*Table 7. Retail cost of methane SCP for different cost scenarios in U.S. dollars per kilogram of dry SCP product.*

Scenario	GCFS conditions (6 years plant lifetime, 24/7 construction)		Regular conditions (20 years plant lifetime, regular construction)	
Energy and feedstock cost	Free NG, low electricity	High	Free NG, low electricity	High
Wholesale price (USD/kg)	\$1.51	\$2.58	\$0.77	\$1.84
Retail cost (USD/kg)	\$3.02	\$5.16	\$1.54	\$3.68

## 4 Discussion

The product retail cost for GCFS conditions does not significantly increase when comparing the scenarios of low versus high operating cost, the difference being US\$3-5/kg dry. The difference in retail cost between regular and GCFS conditions is estimated at approximately \$2/kg. In an ASRS the sunlight levels could be reduced for longer than 6 years, but even with this longer delay before sunlight recovered to current levels, factories built later would still have fewer years of operation during which SCP products could command higher prices due to high demand. There would be lower demand for the SCP product after agricultural productivity recovers, although there could be some remaining opportunity for these plants to operate. The cost analysis is likely conservative for factories operating in the first year of the ASRS. At this price, it is expected that around 90% of the global population would be able to afford the SCP product for all calories with current incomes (Denkenberger et al., 2019).

An important caveat regarding the cost estimations is that the financial assumptions used are common during business as usual, but the financial conditions of an ASRS are complex and outside of the scope of this work. There is considerable uncertainty; although governments gave interest-free loans during the COVID-19 pandemic, raising capital during a GCFS could conversely become far more difficult in the financial ecosystem. Further research is needed on market equilibrium during an ASRS for more precise price estimations of resilient foods.

As shown in Figure 3, it is conceivable that most or all of the methane required to fulfill the global protein requirements could be sourced exclusively from a combination of biogas and natural gas associated with oil which is currently being flared or reinjected. This means that methane SCP production may require redirecting little or no additional natural gas production capacity from current or future uses in order to fulfill this goal. Locations with abundance of flared or vented natural gas should be prioritized for lower production costs. Additionally, these would not require

redirecting the current natural gas production from its use as fuel, which would be a significant advantage during a nuclear winter. Using the natural gas feedstock as an energy source for fulfilling the thermal and electricity energy requirements of the plant could be more convenient than using other sources, but doing so would reduce the amount of feedstock available for further ramp-up. Additionally, further research on the availability of biogas during an ASRS would help provide more insight on the share of natural gas production capacity that would have to be redirected. Biogas produced from decaying plant biomass killed by an ASRS could be significant. Biogas could likely more efficiently be leveraged by co-cultivation of the methane-oxidizing bacteria with hydrogen-oxidizing bacteria (Kerckhof et al., 2021).

Methane SCP is generally faster to ramp up compared to other industrial solutions for resilient food production in ASRS, such as new construction of lignocellulosic sugar plants (Throup et al., 2022) or H<sub>2</sub> SCP plants (García Martínez et al., 2021b). Other non-industrial, low-tech resilient food solutions such as tropical greenhouses (Alvarado et al., 2020) and seaweed farming in the ocean (Mill et al., 2019) are expected to scale up production faster. However, the high protein content and quality of the methane SCP product far surpasses that of these faster scaling solutions, making it valuable as a protein supplementation food during a GCFS. For this reason, a recommendation would be to limit the ramp-up of methane SCP to the production capacity required to fulfill global protein requirements at most, while the rest of the calories and nutrients are fulfilled by faster scaling resilient food solutions that the SCP nutritionally complements (Pham et al., 2022).

Regardless of the ramp-up speed of methane SCP, it would be a great advantage to have multiple factories built and operating before the strike of a GCFS. This would imply a head-start in terms of food production, equivalent to shifting the ramp-up curves upward. If the factories had originally been built for producing animal feed, adding a nucleic acid removal treatment would allow obtaining valuable human food early in the catastrophe. For each plant (of the reference size) available at the onset of an ASRS, an estimated maximum of 620,000 people could meet their entire caloric needs with the SCP product; an estimated maximum of 2.6 million people could meet their minimum protein requirements through the product. Some potential interventions to expedite the ramp-up of SCP technology that were previously identified apply for the case of methane SCP. A coordinated response plan for materials and labor deployment to a collection of pre approved sites would accelerate the initial response to the GCFS. A readily available, generalist methane SCP plant front-end engineering design package, whether produced by industry or academia, could serve as a design basis for the new plants, similar to how (Humbird et al., 2011) serves as a benchmark case study for the biochemical ethanol production process. In addition, once the sector has reached maturity industry experts could create a guide on building and operating the plants, apart from sharing lessons on how to successfully reach and maintain production at scale (García Martínez et al., 2021b). Studies have estimated that effective work on resilience to GCFS is highly cost-effective for the long-term (Denkenberger et al., 2022) as well as for the current generation (Denkenberger and Pearce, 2016, 2018a).

As shown in Figure 5, the protein content of the SCP product significantly influences the speed at which the global protein requirements can be fulfilled. If feasible, research on increasing the protein content would be of important use to this end, be it via optimization of operational parameters such as the methane to nitrogen ratio (Valverde-Pérez et al., 2020), use of thermophilic methanotrophs for reduced cooling requirements (El Abbadi et al., 2022), genetic modification of the microbes, or other means. These types of optimization, if performed prior to a GCFS, would increase the potential of SCP as a resilient food.

The basis of the ramp up model is the assumption that only the capital budget of chemical and related industries (489 billion USD) could be leveraged for construction of SCP factories, roughly accounting in this way for uncertainties in availability of construction materials, construction of chemical equipment and retraining of qualified labor. A previous order-of-magnitude estimate based on growth-rate values suggested that 100% of global human caloric requirements could be fulfilled by methane SCP at around 6 months (Denkenberger and Pearce, 2015), in comparison to the 10 years estimated in Section 3.3. However, the current estimate is considered to be conservative. If the potential of the global construction industry to provide these resources could be effectively leveraged in its entirety, methane SCP could potentially fulfill the entire global caloric requirements of humanity in the time it would take to build a single wave of factories using 24/7 construction, or just over a year. This is because the annual construction expenditure is estimated at 12 trillion USD (de Best, 2021), surpassing the capital requirements for fast construction of methane SCP at 6.1 trillion USD for fulfilling caloric requirements.

Even with the current conservative estimate, there seem to be no significant impediments to securing the global protein supply using methane SCP within a nuclear winter period. Nonetheless, the time this would take is still longer than the 3-6 months period that global food reserves are expected to last (Denkenberger et al., 2019), and would require unprecedented international cooperation. More research is needed on these labor, equipment construction and coordination constraints, so that the degree to which they would hinder the ramp up of methane SCP production (or production of other resilient foods for GCFS) is better understood. Similar to how a wide array of distributed production occurred to overcome supply shortages during the COVID-19 pandemic, it is expected that distributed production of SCP technologies would create greater resiliency in the food system. Similarly to the calls for open hardware to prevent medical shortages (Chagas et al., 2020; Pearce, 2020b, 2020a) the development of open source scalable components of the small-scale SCP production systems is possible. Some of the units are amenable to additive manufacturing and thus distributed open source production. There has already been substantial progress made on various heat exchangers (Arie et al., 2017; Denkenberger and Pearce, 2018b; Tiwari et al., 2019). In addition, there are open source desktop centrifuges that could be scaled for this application (WareJoncas et al., 2018; Sule et al., 2019). Additional work is needed to develop low-cost open source and easily manufactured flash evaporators, spray dryers, cell homogenizers, gas compressors and appropriate bioreactors with high gas utilization.

Low-tech alternatives for downstream processing in small scale SCP production could be settling trays in a refrigerator instead of a centrifuge and boiling the resulting concentrate instead of using a cell homogenizer and spray dryer. Small scale production could use air as O<sub>2</sub> source to obviate the need for air separation, at the cost of reduced gas utilization. Though small scale production of methane SCP would be less efficient, it could have the feature of the waste heat heating the house (particularly advantageous in the case of nuclear winter). There would not be additional retailing costs in contrast to large scale production. Safety issues, and the impact of fugitive methane emissions from small, less controlled "home" or "community" SCP production facilities would have to be assessed. We consider large, industrial SCP production plants more realistic than small-scale efforts, as the process requires proper control. Supplying strains and nutrients for local small-scale productions also seems hard to achieve.

Repurposing existing infrastructure for SCP production could expedite GCFS response, and has already been studied for sugar production in the ASRS context (Throup et al., 2022). Similar to the methodology used in the sugar work, a unit to unit comparison of the methane SCP production

process was made with several different industries, including breweries, instant coffee, milk powder, whey protein, washing powder, cheese powder, fertilizers and biorefineries, showing some overlap. Whey protein factories already have units present for centrifugation, spray drying, packaging and storage of powdered product, but the global whey production capacity is insignificant compared to that required to fulfill global protein requirements (~1% of 0.24-0.39 Gt SCP/year required). On the other hand, the global production capacity of cheese powder and fertilizer factories together amount to a larger tonnage than would be required to fulfill protein requirements via SCP production (0.093 and 0.317 Gt respectively (FAO, 2017; GIR, 2020)). If the drying process present in those factories could be leveraged in combination with the utilities, packing, power, steam, buildings and facilities, savings upwards of 40% of the total capital cost could potentially be obtained based on the Unibio reference plant (Jorgensen, 2011). The value is in accordance with a previously published analysis on repurposing paper mills to biorefineries, which similarly predicts capital savings of 40%, mostly from leveraging existing buildings and service facilities (Martinkus and Wolcott, 2017). However, the very specific nature of the equipment required for SCP production could make it impracticable to repurpose existing units, particularly some of the costliest ones such as reactors, centrifuges and spray dryers. Future research will study the potential of repurposing existing infrastructure for SCP production in depth given the significance of expediting GCFS response.

This needed future research can be viewed as a relatively high priority for GCFS mitigation and preparedness funding, but could also be viewed as a means of insurance for conventional methane infrastructure. As the science behind climate change becomes more granular, risks and liabilities for companies responsible for GHG emissions increase (Allen, 2003; Kunreuther and Michel-Kerjan, 2007; Faure and Peeters, 2011; Heidari and Pearce, 2016) and there is a mounting pressure to mitigate climate liability risk. Although natural gas is less carbon intensive than coal, natural gas pipelines have recently been identified as high priority targets for strategic lawsuits to act as GHG emissions bottlenecks (Pascaris and Pearce, 2020). Natural gas companies could argue that their infrastructure could provide security during GCFS events that impact food systems as they make the transition to hydrogen carriers in a green economy if the future work outlined here to make SCP viable and easily scaled is funded and deployed globally.

## 5 Conclusions

The estimated capital cost for a methane SCP facility for human food production is \$4,800/tpa when built using 24/7 construction to quickly produce protein-rich food during a global catastrophic food shock. The expected retail cost of the product would be in the range of \$3-5/kg (dry), depending on the cost of the natural gas feedstock and electricity.

The ramp up time of methane SCP is estimated at 2.5-4.5 years for fulfilling the protein requirements of the global population, within the duration of a nuclear winter scenario. No further scaling is recommended given the presence of faster scaling resilient foods. 7-11% of global protein requirements could be fulfilled at the end of the first year. Any facilities constructed prior to an abrupt sunlight reduction scenario would imply an important head start in food production capacity, given the sudden nature of the catastrophe and limited window for adaptation.

No significant resource bottlenecks have been found regarding the inputs of methane SCP required to fulfill global protein requirements. Potential bottlenecks regarding availability of construction materials, construction of chemical equipment and retraining of qualified labor were considered limiting. If these were not significant, protein requirements could potentially be fulfilled in little over a year. More research is needed on the degree to which these bottlenecks would affect SCP ramp-up speed.

Other important future research could include the repurposing potential of existing facilities, the economic conditions and biogas potential for methane production during an abrupt sunlight reduction scenario, or the creation of open source designs for small scale production. Recommended interventions include stimulating SCP production during business as usual, creation of a coordinated response plan for materials and construction and development of open generalist SCP front-end engineering design packages and guides.

In conclusion, methane SCP has significant potential to prevent global protein starvation during a global catastrophic food shock at a price affordable by most of the world's population.

## **Acknowledgements**

Thanks to Dr. Silvio Matassa for his help and insights. Thanks to Megan Jamer for editing.

## **Funding**

This research was supported by the Alliance to Feed the Earth in Disasters (ALLFED). Dr. Joshua Pearce is funded by the Thompson Endowment.

## **Conflict of interest**

Dr. Maximilian Lackner is employed by Circe Biotechnologie GmbH. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## References

- Acosta, N., Sakarika, M., Kerckhof, F.-M., Law, C. K. Y., De Vrieze, J., and Rabaey, K. (2020). Microbial protein production from methane via electrochemical biogas upgrading. *Chem. Eng. J.* 391, 123625. doi: 10.1016/j.cej.2019.123625.
- Adjei, A. A., Yamamoto, S., and Kulkarni, A. (1995). Nucleic Acids and/or Their Components: A Possible Role in Immune Function. *J. Nutr. Sci. Vitaminol. (Tokyo)* 41, 1–16. doi: 10.3177/jnsv.41.1.
- Ahn, S., and Norwood, F. B. (2020). Measuring Food Insecurity during the COVID-19 Pandemic of Spring 2020. *Appl. Econ. Perspect. Policy.* doi: 10.1002/aepp.13069.
- Allen, M. (2003). Liability for climate change. *Nature* 421, 891–892. doi: 10.1038/421891a.
- Alptekin, G. (2017). Low Cost Air Separation Process for Gasification Applications. Available at: [https://netl.doe.gov/sites/default/files/event-proceedings/2017/crosscutting/20170320-Track-C/20170320\\_1330C\\_Presentation\\_FE0026142\\_TDAResearch.pdf](https://netl.doe.gov/sites/default/files/event-proceedings/2017/crosscutting/20170320-Track-C/20170320_1330C_Presentation_FE0026142_TDAResearch.pdf) [Accessed July 24, 2020].
- Alvarado, K. A., García Martínez, J. B., Matassa, S., Egbejimba, J., and Denkenberger, D. (2021). Food in space from hydrogen-oxidizing bacteria. *Acta Astronaut.* 180, 260–265. doi: 10.1016/j.actaastro.2020.12.009.
- Alvarado, K. A., Mill, A., Pearce, J. M., Vocaet, A., and Denkenberger, D. (2020). Scaling of greenhouse crop production in low sunlight scenarios. *Sci. Total Environ.* 707, 136012. doi: 10.1016/j.scitotenv.2019.136012.
- Anderson, C., and Solomons, G. L. (1984). "Primary metabolism and biomass production from Fusarium," in *The Applied Mycology of Fusarium*, eds. M. O. Moss and J. E. Smith (Cambridge: Cambridge University Press). Available at: <https://www.osti.gov/etdeweb/biblio/5049931> [Accessed February 19, 2021].
- Aneke, M., and Wang, M. (2015). Potential for improving the energy efficiency of cryogenic air separation unit (ASU) using binary heat recovery cycles. *Appl. Therm. Eng.* 81, 223–231. doi: 10.1016/j.applthermaleng.2015.02.034.
- Arie, M. A., Shooshtari, A. H., Tiwari, R., Dessiatoun, S. V., Ohadi, M. M., and Pearce, J. M. (2017). Experimental characterization of heat transfer in an additively manufactured polymer heat exchanger. *Appl. Therm. Eng.* 113, 575–584. doi: 10.1016/j.applthermaleng.2016.11.030.
- Attanasi, E. D., and Freeman, P. A. (2013). Role of Stranded Gas in Increasing Global Gas Supplies. Reston, Virginia: U.S. Geological Survey Available at: <https://pubs.usgs.gov/of/2013/1044/> [Accessed September 25, 2020].
- Babi, D. K., and Price, J. (2010). Single Cell Protein from Landfill Gas. in (Lyngby, Denmark: Technical University of Denmark). doi: 10.4122/1.1000000046.
- Bailey, R., Benton, T. G., Challinor, A., Elliott, J., Gustafson, D., Hiller, B., et al. (2015). Extreme weather and resilience of the global food system: Final Project Report from the UK-US Taskforce on Extreme Weather and Global Food System Resilience. London: UK: The Global Food Security Programme Available at: [https://www.stat.berkeley.edu/~aldous/157/Papers/extreme\\_weather\\_resilience.pdf](https://www.stat.berkeley.edu/~aldous/157/Papers/extreme_weather_resilience.pdf).
- Baker, C., and McKenzie, K. (2005). Energy consumption of industrial spray dryers. *Dry. Technol.* 23, 365–386. doi: 10.1081/drt-200047665.
- Beasley, D. (2020). WFP Chief warns of hunger pandemic as COVID-19 spreads (Statement to UN Security Council) | World Food Programme. Available at: <https://www.wfp.org/news/wfp-chief-warns-hunger-pandemic-covid-19-spreads-statement-un-security-council> [Accessed May 18, 2020].

- Betti, F., and Heinzmann, T. (2020). COVID-19: How companies are changing track to join the fight. *World Econ. Forum*. Available at: <https://www.weforum.org/agenda/2020/03/from-perfume-to-hand-sanitiser-tvs-to-face-masks-how-companies-are-changing-track-to-fight-covid-19/> [Accessed May 27, 2020].
- Bostrom, N. (2013). Existential Risk Prevention as Global Priority. *Glob. Policy* 4, 15–31. doi: 10.1111/1758-5899.12002.
- Bostrom, N., and Cirkovic, M. M. (2011). *Global catastrophic risks*. Oxford: Oxford University Press.
- Brosemer, K., Schelly, C., Gagnon, V., Arola, K. L., Pearce, J. M., Bessette, D., et al. (2020). The energy crises revealed by COVID: Intersections of Indigeneity, inequity, and health. *Energy Res. Soc. Sci.* 68, 101661. doi: 10.1016/j.erss.2020.101661.
- Burns, S. (2015). Power Costs in the Production of Primary Aluminum. *Steel Alum. Copp. Stainl. Rare Earth Met. Prices Forecast. Met.* Available at: <https://agmetalminer.com/2015/11/24/power-costs-the-production-primary-aluminum/> [Accessed May 20, 2020].
- Calysta Inc. (2020). Adisseo and Calysta establish a Joint-Venture to commercialize FeedKind®. Available at: <http://www.feedkind.com/adisseo-calysta-establish-joint-venture-commercialize-feedkind/> [Accessed May 23, 2020].
- Chagas, A. M., Molloy, J. C., Prieto-Godino, L. L., and Baden, T. (2020). Leveraging open hardware to alleviate the burden of COVID-19 on global health systems. *PLOS Biol.* 18, e3000730. doi: 10.1371/journal.pbio.3000730.
- Christi, M. Y. (1989). *Airlift bioreactors*. London: Elsevier Applied Science.
- Circe Biotech (2021). Protein: feed & food. Available at: <https://www.circe.at/products/protein> [Accessed February 16, 2022].
- Cirkovic, M. M. (2008). "Observation selection effects and global catastrophic risks," in *Global catastrophic risks* (Oxford University Press Oxford), 120–145.
- Claassens, N. J., Sousa, D. Z., dos Santos, V. A. P. M., de Vos, W. M., and van der Oost, J. (2016). Harnessing the power of microbial autotrophy. *Nat. Rev. Microbiol.* 14, 692–706. doi: 10.1038/nrmicro.2016.130.
- Cotton-Barratt, O., Daniel, M., and Sandberg, A. (2020). Defence in Depth Against Human Extinction: Prevention, Response, Resilience, and Why They All Matter. *Glob. Policy* 11, 271–282. doi: 10.1111/1758-5899.12786.
- Coupe, J., Bardeen, C. G., Robock, A., and Toon, O. B. (2019). Nuclear winter responses to nuclear war between the United States and Russia in the whole atmosphere community climate model version 4 and the Goddard Institute for Space Studies ModelE. *J. Geophys. Res. Atmospheres* 124, 8522–8543. doi: <https://doi.org/10.1029/2019JD030509>.
- Crumbley, A. M., and Gonzalez, R. (2018). "Cracking 'Economies of Scale': Biomanufacturing on Methane-Rich Feedstock," in *Methane Biocatalysis: Paving the Way to Sustainability*, eds. M. G. Kalyuzhnaya and X.-H. Xing (Cham: Springer International Publishing), 271–292. doi: 10.1007/978-3-319-74866-5\_17.
- Cumberlege, T., Blenkinsopp, T., and Clark, J. (2016). Assessment of environmental impact of FeedKind protein. 26. doi: <https://prod-drupal-files.storage.googleapis.com/documents/resource/public/Assessment%20of%20environmental%20footprint%20of%20FeedKind%20protein%20-%20REPORT.pdf>.
- Damodaran, A. (2020). Global Capital Expenditures, Acquisitions and R&D and Sales/Invested Capital Ratios. Available at: <http://www.stern.nyu.edu/~adamodar/pc/datasets/capexGlobal.xls> [Accessed September 3, 2020].

- de Best, R. (2021). Construction industry spending globally. *Stat. Constr. Ind. Spend. Worldw. 2014–2019 Forecasts 2020–2035*. Available at: <https://www.statista.com/statistics/788128/construction-spending-worldwide/> [Accessed February 19, 2021].
- Denkenberger, D. C., Cole, D. D., Abdelkhaliq, M., Griswold, M., Hundley, A. B., and Pearce, J. M. (2017). Feeding everyone if the sun is obscured and industry is disabled. *Int. J. Disaster Risk Reduct.* 21, 284–290. doi: 10.1016/j.ijdrr.2016.12.018.
- Denkenberger, D. C., and Pearce, J. M. (2015). Feeding everyone: Solving the food crisis in event of global catastrophes that kill crops or obscure the sun. *Futures* 72, 57–68. doi: <https://doi.org/10.1016/j.futures.2014.11.008>.
- Denkenberger, D. C., and Pearce, J. M. (2016). Cost-effectiveness of interventions for alternate food to address agricultural catastrophes globally. *Int. J. Disaster Risk Sci.* 7, 205–215. doi: 10.1007/s13753-016-0097-2.
- Denkenberger, D. C., and Pearce, J. M. (2018a). Cost-effectiveness of interventions for alternate food in the United States to address agricultural catastrophes. *Int. J. Disaster Risk Reduct.* 27, 278–289. doi: <https://doi.org/10.1016/j.ijdrr.2017.10.014>.
- Denkenberger, D., and Pearce, J. M. (2014). *Feeding everyone no matter what: Managing food security after global catastrophe*. Cambridge, Massachusetts: Academic Press.
- Denkenberger, D., Pearce, J., Taylor, A. R., and Black, R. (2019). Food without sun: Price and life-saving potential. *Foresight* 21, 118–129. doi: 10.1108/FS-04-2018-0041.
- Denkenberger, D., Sandberg, A., Tieman, R. J., and Pearce, J. M. (2021). Long-term cost-effectiveness of interventions for loss of electricity/industry compared to artificial general intelligence safety. *Eur. J. Futur. Res.* 9, 11. doi: 10.1186/s40309-021-00178-z.
- Denkenberger, D., Sandberg, A., Tieman, R. J., and Pearce, J. M. (2022). Long term cost-effectiveness of resilient foods for global catastrophes compared to artificial general intelligence safety. *Int. J. Disaster Risk Reduct.* 73, 102798. doi: 10.1016/j.ijdrr.2022.102798.
- Denkenberger, and Pearce, J. M. (2018b). Design Optimization of Polymer Heat Exchanger for Automated Household-Scale Solar Water Pasteurizer. *Designs* 2, 11. doi: 10.3390/designs2020011.
- Duchene, L. (2016). Bridging the omega-3 gap with methane, microalgae. *Glob. Seaf. Alliance*. Available at: <https://www.globalseafood.org/advocate/bridging-the-omega-3-gap-with-methane-microalgae/> [Accessed January 14, 2022].
- EIA (2019). U.S. Energy Information Administration: International Energy Statistics database. Available at: <https://www.eia.gov/international/data/world> [Accessed July 22, 2020].
- EIA (2020a). Average retail price of electricity. *US Energy Inf. Adm.* Available at: <https://www.eia.gov/electricity/data/browser/#/topic/7?agg=2,0,1&geo=g&freq=M> [Accessed May 20, 2020].
- EIA (2020b). Coal Markets. *US Energy Inf. Adm.* Available at: <https://www.eia.gov/coal/markets/> [Accessed May 20, 2020].
- El Abbadi, S. H., Sherwin, E. D., Brandt, A. R., Luby, S. P., and Criddle, C. S. (2022). Displacing fishmeal with protein derived from stranded methane. *Nat. Sustain.* 5, 47–56. doi: 10.1038/s41893-021-00796-2.
- Elvidge, C. D., Zhizhin, M., Baugh, K., Hsu, F.-C., and Ghosh, T. (2016). Methods for global survey of natural gas flaring from visible infrared imaging radiometer suite data. *Energies* 9, 14. doi: <https://doi.org/10.3390/en9010014>.
- Enbridge Gas Inc. (2001). Chemical Composition of Natural Gas. Available at: <https://www.uniongas.com/about-us/about-natural-gas/chemical-composition-of-natural-g>

- as [Accessed August 27, 2020].
- Engineering ToolBox (2005). Fuel Gases Heating Values. Available at:  
[https://www.engineeringtoolbox.com/heating-values-fuel-gases-d\\_823.html](https://www.engineeringtoolbox.com/heating-values-fuel-gases-d_823.html) [Accessed August 27, 2020].
- Eurostat (2019). Development of electricity prices for non-household consumers, EU-28 and EA. Eurostat. Available at:  
[https://ec.europa.eu/eurostat/statistics-explained/index.php?title=File:Development\\_of\\_electricity\\_prices\\_for\\_non-household\\_consumers,\\_EU-28\\_and\\_EA,\\_2008-2019\\_\(EUR\\_per\\_kWh\).png](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=File:Development_of_electricity_prices_for_non-household_consumers,_EU-28_and_EA,_2008-2019_(EUR_per_kWh).png) [Accessed May 20, 2020].
- FAO (2017). World Fertilizer Trends and Outlook to 2020. doi: <http://www.fao.org/3/a-i6895e.pdf>.
- Faure, M., and Peeters, M. (2011). *Climate change liability*. Cheltenham: Edward Elgar Publishing.
- Finnigan, T. J. A., Wall, B. T., Wilde, P. J., Stephens, F. B., Taylor, S. L., and Freedman, M. R. (2019). Mycoprotein: The Future of Nutritious Nonmeat Protein, a Symposium Review. *Curr. Dev. Nutr.* 3. doi: [10.1093/cdn/nzz021](https://doi.org/10.1093/cdn/nzz021).
- Flanagan, S. (2022). Calysta appoints Chief Science and Sustainability Officer. Calysta. Available at:  
<https://calysta.com/calysta-appoints-chief-science-and-sustainability-officer/> [Accessed February 16, 2022].
- FMI (2019). Industrial Oxygen: Meeting Alternative Energy Demands. Dublin: Future Market Insights.
- García Martínez, J. B., Alvarado, K. A., and Denkenberger, D. C. (2022). Synthetic fat from petroleum as a resilient food for global catastrophes: Preliminary techno-economic assessment and technology roadmap. *Chem. Eng. Res. Des.* 177, 255–272. doi: [10.1016/j.cherd.2021.10.017](https://doi.org/10.1016/j.cherd.2021.10.017).
- García Martínez, J. B., Brown, M. M., Christodoulou, X., Alvarado, K. A., and Denkenberger, D. C. (2021a). Potential of microbial electrosynthesis for contributing to food production using CO<sub>2</sub> during global agriculture-inhibiting disasters. *Clean. Eng. Technol.* 4. doi: [10.1016/j.clet.2021.100139](https://doi.org/10.1016/j.clet.2021.100139).
- García Martínez, J. B., Egbejimba, J., Throup, J., Matassa, S., Pearce, J. M., and Denkenberger, D. C. (2021b). Potential of microbial protein from hydrogen for preventing mass starvation in catastrophic scenarios. *Sustain. Prod. Consum.* 25, 234–247. doi: <https://doi.org/10.1016/j.spc.2020.08.011>.
- GIR (2020). Global Cheese Powder Market Size, Forecast Report 2025. Hong Kong: Global Info Research Available at:  
<https://www.decisiondatabases.com/ip/21253-cheese-powder-market-analysis-report> [Accessed September 16, 2020].
- Glencross, B. D., Huyben, D., and Schrama, J. W. (2020). The Application of Single-Cell Ingredients in Aquaculture Feeds—A Review. *Fishes* 5, 22. doi: [10.3390/fishes5030022](https://doi.org/10.3390/fishes5030022).
- Hanna, A. S., Chang, C.-K., Lackney, J. A., and Sullivan, K. T. (2007). Impact of Overmanning on Mechanical and Sheet Metal Labor Productivity. *J. Constr. Eng. Manag.* 133, 22–28. doi: [10.1061/\(ASCE\)0733-9364\(2007\)133:1\(22\)](https://doi.org/10.1061/(ASCE)0733-9364(2007)133:1(22)).
- Heidari, N., and Pearce, J. M. (2016). A review of greenhouse gas emission liabilities as the value of renewable energy for mitigating lawsuits for climate change related damages. *Renew. Sustain. Energy Rev.* 55, 899–908. doi: [10.1016/j.rser.2015.11.025](https://doi.org/10.1016/j.rser.2015.11.025).
- Humbird, D., Davis, R., Tao, L., Kinchin, C., Hsu, D., Aden, A., et al. (2011). Process design and economics for biochemical conversion of lignocellulosic biomass to ethanol: dilute-acid pretreatment and enzymatic hydrolysis of corn stover. National Renewable Energy Lab.(NREL), Golden, CO (United States) Available at:  
<https://www.nrel.gov/docs/fy11osti/47764.pdf>.
- IEA (2019). Natural Gas Information: Overview. Nat. Gas Inf. Overv. Available at:

- <https://www.iea.org/reports/natural-gas-information-overview>.
- Jain, S. (2019). Global Potential of Biogas. doi:  
<https://www.worldbiogasassociation.org/global-potential-of-biogas/>.
- Jenkins, G. (1988). "SCP — The BP Protein Process," in *Resources and Applications of Biotechnology: The New Wave*, ed. R. Greenshields (London: Palgrave Macmillan UK), 141–149. doi: 10.1007/978-1-349-09574-2\_15.
- Jensen, M. (2021). Unibio solutions for UFMSW valorisation from Valuewaste. doi:  
<https://hoopproject.eu/wp-content/uploads/2021/05/ValueWaste-Project-PPT-Presentation-26052021.pptx-pdf.pdf>.
- Jones, S., Karpol, A., Friedman, S., Maru, B., and Tracy, B. (2020). Recent advances in single cell protein use as a feed ingredient in aquaculture. *Curr. Opin. Biotechnol.* 61, 189–197. doi: 10.1016/j.copbio.2019.12.026.
- Jorgensen, J. B. (2011). Exploiting Microorganisms for Animal Feed Production. in (The 7th Malaysia International Agro-Bio Business Conference : What Next Agriculture II, Kuala Lumpur, Malaysia). doi: <https://core.ac.uk/download/pdf/13790016.pdf>.
- Kerckhof, F.-M., Sakarika, M., Van Giel, M., Muys, M., Vermeir, P., De Vrieze, J., et al. (2021). From Biogas and Hydrogen to Microbial Protein Through Co-Cultivation of Methane and Hydrogen Oxidizing Bacteria. *Front. Bioeng. Biotechnol.* 9, 771. doi: 10.3389/fbioe.2021.733753.
- Khoshnevisan, B., He, L., Xu, M., Valverde-Pérez, B., Sillman, J., Mitraka, G.-C., et al. (2022). From renewable energy to sustainable protein sources: Advancement, challenges, and future roadmaps. *Renew. Sustain. Energy Rev.* 157, 112041. doi: 10.1016/j.rser.2021.112041.
- Khoshnevisan, B., Tsapekos, P., Zhang, Y., Valverde-Pérez, B., and Angelidaki, I. (2019). Urban biowaste valorization by coupling anaerobic digestion and single cell protein production. *Bioresour. Technol.* 290, 121743. doi: 10.1016/j.biortech.2019.121743.
- Kreutz, T., Williams, R., Consonni, S., and Chiesa, P. (2005). Co-production of hydrogen, electricity and CO<sub>2</sub> from coal with commercially ready technology. Part B: Economic analysis. *Int. J. Hydrol. Energy* 30, 769–784. doi: 10.1016/j.ijhydene.2004.08.001.
- Kunreuther, H. C., and Michel-Kerjan, E. O. (2007). Climate change, insurability of large-scale disasters and the emerging liability challenge. in 6. (Cambridge, Massachusetts: The University of Pennsylvania Law Review). doi: 10.3386/w12821.
- Laborde, D., Martin, W., and Vos, R. (2020). Poverty and food insecurity could grow dramatically as COVID-19 spreads. *Int. Food Policy Res. Inst. IFPRI Wash. DC*. doi:  
<https://www.ifpri.org/publication/poverty-and-food-insecurity-could-grow-dramatically-covid-19-spreads>.
- Lane, J. (2018). Construction complete at Unibio/Protelux commercial scale methane to protein plant : Biofuels Digest. Available at:  
<http://www.biofuelsdigest.com/bdigest/2018/09/13/construction-complete-at-unibio-protelux-commercial-scale-methane-to-protein-plant/> [Accessed May 23, 2020].
- Lee, A. (2021). Barriers to adopting alternative protein are decreasing, post-Covid-19: String Bio co-founder Ezhil Subbian. Eco-Bus. Available at:  
<https://www.eco-business.com/news/barriers-to-adopting-alternative-protein-are-decreasing-post-covid-19-string-bio-co-founder-ezhil-subbian/> [Accessed September 13, 2021].
- Leger, D., Matassa, S., Noor, E., Shepon, A., Milo, R., and Bar-Even, A. (2021). Photovoltaic-driven microbial protein production can use land and sunlight more efficiently than conventional crops. *Proc. Natl. Acad. Sci.* 118. doi: 10.1073/pnas.2015025118.
- Levett, I., Birkett, G., Davies, N., Bell, A., Langford, A., Laycock, B., et al. (2016). Techno-economic assessment of poly-3-hydroxybutyrate (PHB) production from methane—The case for thermophilic bioprocessing. *J. Environ. Chem. Eng.* 4, 3724–3733. doi:

10.1016/j.jece.2016.07.033.

- Manero, A., Smith, P., Koontz, A., Dombrowski, M., Sparkman, J., Courbin, D., et al. (2020). Leveraging 3D Printing Capacity in Times of Crisis: Recommendations for COVID-19 Distributed Manufacturing for Medical Equipment Rapid Response. *Int. J. Environ. Res. Public. Health* 17. doi: 10.3390/ijerph17134634.
- Manheim, D., and Denkenberger, D. (2020). Review of potential high-leverage and inexpensive mitigations for reducing risk in epidemics and pandemics. *J. Glob. Health Rep.* 4, e2020010. doi: 10.29392/001c.12530.
- Markets Insider (2020a). Coal PRICE Today | Coal Spot Price Chart. [markets.businessinsider.com/commodities/coal-price](https://markets.businessinsider.com/commodities/coal-price) [Accessed May 20, 2020].
- Markets Insider (2020b). Natural Gas PRICE Today. *Nat. Gas PRICE Today*. Available at: <https://markets.businessinsider.com/commodities/natural-gas-price> [Accessed August 28, 2020].
- Marlow Foods Ltd. (2001). GRAS NOTIFICATION for MYCOPROTEIN. Available at: <http://wayback.archive-it.org/7993/20171031053437/https://www.fda.gov/downloads/Food/IngredientsPackagingLabeling/GRAS/NoticeInventory/UCM266876.pdf> [Accessed July 30, 2020].
- Martin, J., Burrows, T. K., and Pegg, I. (2006). Predicting Construction Duration of Building Projects. in (Munich). doi: [https://fig.net/resources/proceedings/fig\\_proceedings/fig2006/papers/ts28/ts28\\_02\\_martin\\_eta\\_0831.pdf](https://fig.net/resources/proceedings/fig_proceedings/fig2006/papers/ts28/ts28_02_martin_eta_0831.pdf).
- Martinkus, N., and Wolcott, M. (2017). A framework for quantitatively assessing the repurpose potential of existing industrial facilities as a biorefinery. *Biofuels Bioprod. Biorefining* 11, 295–306. doi: <https://doi.org/10.1002/bbbb.1742>.
- Matassa, S., Batstone, D. J., Hülsen, T., Schnoor, J., and Verstraete, W. (2015). Can direct conversion of used nitrogen to new feed and protein help feed the world? *Environ. Sci. Technol.* 49, 5247–5254. doi: 10.1021/es505432w.
- Matassa, S., Boon, N., Pikaar, I., and Verstraete, W. (2016). Microbial protein: future sustainable food supply route with low environmental footprint. *Microb. Biotechnol.* 9, 568–575. doi: 10.1111/1751-7915.12369.
- Matassa, S., Papirio, S., Pikaar, I., Hülsen, T., Leijenhorst, E., Esposito, G., et al. (2020). Upcycling of biowaste carbon and nutrients in line with consumer confidence: the “full gas” route to single cell protein. *Green Chem.* 22, 4912–4929. doi: 10.1039/DoGCo1382J.
- McCray, B. (2010). How to set retail prices and markups. *Small Biz Surviv.* Available at: <https://smallbizsurvival.com/2010/11/how-to-set-retail-prices-and-markups.html> [Accessed September 22, 2020].
- Mill, A., Harrison, C., James, S., Shah, S., Fist, T., Alvarado, K., et al. (2019). Preventing global famine in case of sun-blocking scenarios: Seaweed as an alternative food source. in doi: [https://www.researchgate.net/publication/337199859\\_Preventing\\_global\\_famine\\_in\\_case\\_of\\_sun-blocking\\_scenarios\\_Seaweed\\_as\\_an\\_alternative\\_food\\_source\\_Key\\_findings](https://www.researchgate.net/publication/337199859_Preventing_global_famine_in_case_of_sun-blocking_scenarios_Seaweed_as_an_alternative_food_source_Key_findings).
- Pascaris, A. S., and Pearce, J. M. (2020). U.S. Greenhouse Gas Emission Bottlenecks: Prioritization of Targets for Climate Liability. *Energies* 13, 3932. doi: 10.3390/en13153932.
- Pearce, J. M. (2020a). A review of open source ventilators for COVID-19 and future pandemics. *F1000Research* 9. doi: 10.12688/f1000research.22942.2.
- Pearce, J. M. (2020b). Distributed Manufacturing of Open Source Medical Hardware for Pandemics. *J. Manuf. Mater. Process.* 4, 49. doi: 10.3390/jmmp4020049.
- Pearce, J. M., Khaksari, M., and Denkenberger, D. (2019). Preliminary Automated Determination of

- Edibility of Alternative Foods: Non-Targeted Screening for Toxins in Red Maple Leaf Concentrate. *Plants* 8, 110. doi: 10.3390/plants8050110.
- Pham, A., García Martínez, J. B., Brynch, V., Stormbjorne, R., Pearce, J. M., and Denkenberger, D. C. (2022). Nutrition in Abrupt Sunlight Reduction Scenarios: Envisioning Feasible Balanced Diets on Resilient Foods. *Nutrients* 14, 492. doi: 10.3390/nu14030492.
- Pikaar, I., Matassa, S., Bodirsky, B. L., Weindl, I., Humpenöder, F., Rabaey, K., et al. (2018). Decoupling livestock from land use through industrial feed production pathways. *Environ. Sci. Technol.* 52, 7351–7359. doi: <https://doi.org/10.1021/acs.est.8b00216>.
- Rajesh, M., Kamalam, B. S., Sharma, P., Verma, V. C., Pandey, A., Dubey, M. K., et al. (2022). Evaluation of a novel methanotroph bacteria meal grown on natural gas as fish meal substitute in rainbow trout, *Oncorhynchus mykiss*. *Aquac. Res.* doi: 10.1111/are.15735.
- Ravindra, P. (2000). Value-added food: Single cell protein. *Biotechnol. Adv.* 18, 459–479. doi: [https://doi.org/10.1016/S0734-9750\(00\)00045-8](https://doi.org/10.1016/S0734-9750(00)00045-8).
- Research and Markets (2020). Ammonia Market - Growth, Trends, and Forecast (2020 - 2025). Dublin: Research and Markets ltd Available at: <https://www.researchandmarkets.com/reports/4773617/ammonia-market-growth-trends-and-forecast> [Accessed May 14, 2020].
- Ritala, A., Häkkinen, S. T., Toivari, M., and Wiebe, M. G. (2017). Single cell protein—state-of-the-art, industrial landscape and patents 2001–2016. *Front. Microbiol.* 8, 2009. doi: 10.3389/fmicb.2017.02009.
- Rob Smith (2018). These are the world's biggest coal producers. *World Econ. Forum*. Available at: <https://www.weforum.org/agenda/2018/01/these-are-the-worlds-biggest-coal-producers/> [Accessed May 14, 2020].
- Robock, A., Oman, L., and Stenchikov, G. L. (2007). Nuclear winter revisited with a modern climate model and current nuclear arsenals: Still catastrophic consequences. *J. Geophys. Res. Atmospheres* 112. doi: 10.1029/2006JD008235.
- Rosenberry, B. (2019). NovoNutrients: Making Aquatic Feeds from Waste Products. Available at: <https://www.shrimpnews.com/FreeReportsFolder/SpecialReports/NovoNutrientsShrimpFeedsFromWasteProducts.html> [Accessed May 23, 2020].
- Sefton, B. (2018). NovoNutrients: Food from CO<sub>2</sub>. doi: <http://nas-sites.org/dels/files/2018/02/2-2-SEFTON-NovоНutrients-NAS.pdf>.
- Shepon, A., Eshel, G., Noor, E., and Milo, R. (2016). Energy and protein feed-to-food conversion efficiencies in the US and potential food security gains from dietary changes. *Environ. Res. Lett.* 11, 105002. doi: 10.1088/1748-9326/11/10/105002.
- Shokrani, A., Loukaides, E. G., Elias, E., and Lunt, A. J. G. (2020). Exploration of alternative supply chains and distributed manufacturing in response to COVID-19; a case study of medical face shields. *Mater. Des.* 192, 108749. doi: 10.1016/j.matdes.2020.108749.
- Short, W., Packey, D. J., and Holt, T. (1995). A manual for the economic evaluation of energy efficiency and renewable energy technologies. National Renewable Energy Lab., Golden, CO (United States).
- Sillman, J., Nygren, L., Kahiluoto, H., Ruuskanen, V., Tamminen, A., Bajamundi, C., et al. (2019). Bacterial protein for food and feed generated via renewable energy and direct air capture of CO<sub>2</sub>: Can it reduce land and water use? *Glob. Food Secur.* 22, 25–32. doi: 10.1016/j.gfs.2019.09.007.
- Silverman, J. (2020). Single Cell Protein: A sustainable approach to meeting the growing protein demand. *AQUAFEED Adv. Process. Formul.* 7, 10–14.
- Solar Foods (2021). Solein submitted to the European Commission for novel food approval. Sol. Foods Oy. Available at:

- <https://solarfoods.fi/our-news/solein-submitted-to-the-european-commission-for-novel-food-approval/> [Accessed February 17, 2022].
- Sönnichsen, N. (2020). Electricity consumption globally. Statista. Available at: <https://www.statista.com/statistics/280704/world-power-consumption/> [Accessed May 14, 2020].
- Southey, F. (2019). Solar Foods makes protein out of thin air: 'This is the most environmentally friendly food there is.' foodnavigator.com. Available at: <https://www.foodnavigator.com/Article/2019/07/15/Solar-Foods-makes-protein-out-of-thin-air-This-is-the-most-environmentally-friendly-food-there-is> [Accessed May 8, 2020].
- Sule, S. S., Petsiuk, A. L., and Pearce, J. M. (2019). Open Source Completely 3-D Printable Centrifuge. *Instruments* 3, 30. doi: 10.3390/instruments3020030.
- Switzenbaum, M. S., Giraldo-Gomez, E., and Hickey, R. F. (1990). Monitoring of the anaerobic methane fermentation process. *Enzyme Microb. Technol.* 12, 722–730. doi: 10.1016/0141-0229(90)90142-D.
- TEMASEK (2020). The Unlikely Solutions to Close the Protein Gap. Temasek Corp. Website Engl. Available at: <https://www.temasek.com.sg/content/temasek-corporate/en/news-and-views/stories/future/the-next-food-revolution/the-unlikely-solutions-to-close-the-protein-gap> [Accessed September 13, 2021].
- Throup, J., García Martínez, J. B., Bals, B., Cates, J., Pearce, J. M., and Denkenberger, D. C. (2022). Rapid repurposing of pulp and paper mills, biorefineries, and breweries for lignocellulosic sugar production in global food catastrophes. *Food Bioprod. Process.* 131, 22–39. doi: 10.1016/j.fbp.2021.10.012.
- Tiwari, R., Andhare, R. S., Shooshtari, A., and Ohadi, M. (2019). Development of an additive manufacturing-enabled compact manifold microchannel heat exchanger. *Appl. Therm. Eng.* 147, 781–788. doi: 10.1016/j.applthermaleng.2018.10.122.
- Tzachor, A., Richards, C. E., and Holt, L. (2021). Future foods for risk-resilient diets. *Nat. Food* 2, 326–329. doi: 10.1038/s43016-021-00269-x.
- UIG (2006). Universal Industrial Gases Inc. - Optimal System - Plant Selection and Supply System Design. Available at: <http://www.uigi.com/optimalplant.html> [Accessed July 24, 2020].
- Unibio Group (2020a). Chemical Composition of Uniprotein. Unibio. Available at: <https://www.unibio.dk/end-product/chemical-composition-1/> [Accessed May 14, 2020].
- Unibio Group (2020b). Unibio process. Available at: <https://www.unibio.dk/technology/process/> [Accessed September 25, 2020].
- United Nations (2019). World Population Prospects 2019: Highlights. New York Available at: [https://population.un.org/wpp/Publications/Files/WPP2019\\_Highlights.pdf](https://population.un.org/wpp/Publications/Files/WPP2019_Highlights.pdf).
- U.S. Department of Agriculture (2016). How many calories are in one gram of fat, carbohydrate, or protein? | Food and Nutrition Information Center| NAL | USDA. Available at: <https://www.nal.usda.gov/fnic/how-many-calories-are-one-gram-fat-carbohydrate-or-protein> [Accessed July 30, 2020].
- U.S. Food and Drug Administration (2002). GRAS Notices: GRN No. 000091 (Mycoprotein). Available at: <https://www.accessdata.fda.gov/scripts/fdcc/index.cfm?set=GRASNotices&id=91> [Accessed July 30, 2020].
- Valverde-Pérez, B., Xing, W., Zachariae, A. A., Skadborg, M. M., Kjeldgaard, A. F., Palomo, A., et al. (2020). Cultivation of methanotrophic bacteria in a novel bubble-free membrane bioreactor for microbial protein production. *Bioresour. Technol.* 310, 123388. doi: 10.1016/j.biortech.2020.123388.

- Villadsen, J., Nielsen, J., and Lidén, G. (2011). *Bioreaction engineering principles*. New York: Springer Science & Business Media.
- Volova, T. G., and Barashkov, V. A. (2010). Characteristics of proteins synthesized by hydrogen-oxidizing microorganisms. *Appl. Biochem. Microbiol.* 46, 574–579. doi: 10.1134/S0003683810060037.
- WareJoncas, Z., Stewart, C., and Giannini, J. (2018). An Inexpensive, Open-Source Mini-Centrifuge. *Am. Biol. Teach.* 80, 451–456. doi: 10.1525/abt.2018.80.6.451.
- Warner, M. (2019). *Industrial Biotechnology Commercialization Handbook: How to Make Proteins Without Animals and Fuels Or Chemicals Without Crude Oil*. Seattle: Amazon Digital Services LLC.
- World Bank (2018). Global gas flaring and oil production: 1996 - 2018. Available at: <http://pubdocs.worldbank.org/en/182101560185376968/pdf/Global-gas-flaring-and-oil-production-1996-2018.pdf> [Accessed July 16, 2020].
- World Health Organization (2004). Food and nutrition needs in emergencies. doi: <https://www.unhcr.org/uk/45fa745b2.pdf>.
- World Health Organization and United Nations University (2007). *Protein and amino acid requirements in human nutrition*. Geneva Available at: <https://apps.who.int/iris/handle/10665/43411>.
- Worldometers (2020). World Population Clock. *Worldometers*. Available at: <https://www.worldometers.info/world-population/> [Accessed May 28, 2020].
- Xu, M., Zhao, D., Zhu, X., Su, Y., Angelidaki, I., and Zhang, Y. (2021a). Biogas upgrading and valorization to single-cell protein in a bioinorganic electrosynthesis system. *Chem. Eng. J.* 426, 131837. doi: 10.1016/j.cej.2021.131837.
- Xu, M., Zhou, H., Zou, R., Yang, X., Su, Y., Angelidaki, I., et al. (2021b). Beyond the farm: Making edible protein from CO<sub>2</sub> via hybrid bioinorganic electrosynthesis. *One Earth* 4, 868–878. doi: 10.1016/j.oneear.2021.05.007.